



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : A01N 25/30, 57/20	A1	(11) International Publication Number: WO 00/59302 (43) International Publication Date: 12 October 2000 (12.10.00)
<div style="display: flex; justify-content: space-between;"> <div style="width: 45%;"> <p>(21) International Application Number: PCT/GB00/01062</p> <p>(22) International Filing Date: 21 March 2000 (21.03.00)</p> <p>(30) Priority Data: 9907669.7 1 April 1999 (01.04.99) GB</p> <p>(71) Applicant (for all designated States except US): ZENECA LIMITED [GB/GB]; 15 Stanhope Gate, London W1Y 6LN (GB).</p> <p>(72) Inventors; and (75) Inventors/Applicants (for US only): CUTLER, Julia, Lynne [GB/GB]; Jealotts Hill International Research Centre, Bracknell, Berkshire RG42 6ET (GB). BEAN, Michael, John [GB/GB]; Jealotts Hill International Research Centre, Bracknell, Berkshire RG42 6ET (GB). SEVILLE, Antony, George [GB/GB]; Jealotts Hill International Research Centre, Bracknell, Berkshire RG42 6ET (GB).</p> <p>(74) Agents: RICKS, Michael, James et al.; Zeneca Agrochemicals, Intellectual Property Dept., P.O. Box 3538, Jealott's Hill Research Station, Bracknell RG42 6YA (GB).</p> </div> <div style="width: 50%;"> <p>(81) Designated States: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p>Published <i>With international search report.</i></p> </div> </div>		
<p>(54) Title: AGROCHEMICAL COMPOSITION</p> <div style="text-align: center; margin: 20px 0;"> $\begin{array}{c} \text{CR}^{16}\text{R}^{17} - \text{O}- \\ \\ \text{CR}^{18} - (\text{R}^{21})_d - \text{O} - (\text{R}^8\text{O})_b - \text{R}^9 - \text{NR}^{10}\text{R}^{11} \\ \\ \text{CR}^{19}\text{R}^{20} - \text{O} - (\text{R}^{12}\text{O})_c - \text{R}^{13} - \text{NR}^{14}\text{R}^{15} \end{array} \quad (1)$ </div> <p>(57) Abstract</p> <p>An aqueous agrochemical composition comprises an agrochemical active ingredient such as glyphosate, paraquat or fomesafen and an adjuvant of formula (I): $\text{X}-(\text{R}^3\text{O})_a-\text{R}^4-\text{N R}^2\text{R}^1$ and salts thereof, wherein R^1 and R^2 are independently hydrogen or a lower alkyl group or a group $\text{X}-(\text{R}^3\text{O})_a-\text{R}^4-$ wherein X is $-\text{OH}$ or a lower alkoxy group; R^3O is an ethoxy, propoxy or butoxy group or a random or block mixture thereof; R^4 is a linear or branched chain alkylene bridging group containing from 1 to 4 carbon atoms; X is (IA) $-\text{OH}$ or a lower alkyloxy group; or (IB) a group $\text{R}^5\text{R}^6\text{N}-$ or $\text{R}^5\text{R}^6\text{N}-\text{R}^7-$; or (IC) a group (1) wherein a, is from 1 to 400 or if X is of formula IC, the sum of a, b and c is from 3 to 400 and d is 0 or 1.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.


AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

AGROCHEMICAL COMPOSITION

This invention relates to an agrochemical composition and in particular to an agrochemical composition containing an activity-enhancing adjuvant.

5 An agrochemical is generally used with an adjuvant or combination of adjuvants to provide optimum biological activity. Much has been published on the selection of adjuvants to achieve particular effects with individual agrochemicals and classes of agrochemical. In general it has been assumed that activity-enhancement results from surfactant properties of the adjuvant and most such activity-enhancing adjuvants are surfactants in that they contain
10 within the molecule both a hydrophobic portion and a lyophobic portion. We have now found that a class of alkoxyated amines provides excellent activity enhancement when used in an agrochemical formulation. Surprisingly certain members of the class have either no surfactant properties or low surfactant properties. Furthermore, unlike many surfactants used as conventional agrochemical adjuvants, the alkoxyated amines of the present
15 invention generally exhibit an exceptionally low toxicological profile and in particular excellent ecotoxicological characteristics and provide compositions which are especially benign to the environment. It is exceptional to find adjuvants which combine a high level of activity enhancement with low toxicity.

According to the present invention there is provided an agrochemical composition
20 comprising an agrochemical active ingredient and an adjuvant of formula (I) and salts thereof



25 wherein R^1 and R^2 are independently hydrogen or a lower alkyl group or a group $X'-(R^3O)_a-R^4-$ wherein R^3 and R^4 respectively may take any of the values of R^3 and R^4 as hereinafter defined and wherein X' is -OH or a lower alkoxy group containing from 1 to 6 carbon atoms and a is from 0 to 400

R^3O is an ethoxy, propoxy or butoxy group or a random or block mixture thereof,

30 R^4 is a linear or branched chain alkylene bridging group containing from 1 to 4 carbon atoms, X is

(IA) -OH or a lower alkoxy group containing from 1 to 6 carbon atoms

or X is

(IB) a group R^5R^6N- or $R^5R^6N-R^7-$

wherein R^5 and R^6 are independently hydrogen or a lower alkyl group or a group $X'-(R^3O)_a-R^4-$ as hereinbefore defined wherein X' is -OH or a lower alkoxy group

containing from 1 to 6 carbon atoms,

and R^7 is a linear or branched chain alkylene bridging group containing from 1 to 4 carbon atoms

or X is

(IC) a group

$CR^{16}R^{17}-O-$

|

$CR^{18}-(R^{21})_d-O-(R^8O)_b-R^9-NR^{10}R^{11}$

|

$CR^{19}R^{20}-O-(R^{12}O)_c-R^{13}-NR^{14}R^{15}$

wherein R^{10} , R^{11} , R^{14} and R^{15} are independently hydrogen or a lower alkyl group or a group $X'-(R^3O)_a-R^4-$ as hereinbefore defined wherein X' is -OH or a lower alkoxy group containing from 1 to 6 carbon atoms,

and R^8 and R^{12} are independently alkyl groups containing from 1 to 4 carbon atoms and R^9 and R^{13} and R^{21} are independently a linear or branched chain alkylene bridging group containing from 1 to 6 carbon atoms, R^{16} , R^{17} , R^{18} , R^{19} and R^{20} are independently hydrogen or a lower alkyl group and d is 0 or 1

and wherein a, is from 1 to 400 or if X is of formula IC, the sum of a, b and c is from 3 to

400.

A considerable number of salts of the compound of formula (I) suitable for use in agrochemical applications will occur to one skilled in the art. Examples of salts include phosphate, sulphate, carboxylate, acetate, formate and chloride although many other suitable salts will occur to one skilled in the art. Alternatively the compound of formula (I) may form a salt with an acidic agrochemical such as glyphosate. Salts of the compound of formula (I) can also be prepared from acidic surfactants, for example optionally ethoxylated alkyl or alkylene ester derivatives of phosphoric acid or phosphonic acid, or optionally ethoxylated alkyl or alkylene carboxylic acids or sulphonic acids. It will be appreciated that where more than one amine functionality is present in the compound of formula (I) or the

relevant acid has more than one functional group, the option for different salt stoichiometries and mixed salts exists and all such variations are included herein.

As used herein, the term lower alkyl means a linear or branched chain primary or secondary alkyl group containing from 1 to 6 carbon atoms. Preferred lower alkyl groups contain from 1 to 4 carbon atoms, and methyl, ethyl and propyl or isopropyl groups are especially preferred.

R^1 , R^2 , R^5 , R^6 , R^{10} , R^{11} , R^{14} and R^{15} are preferably methyl or hydrogen. Hydrogen is especially preferred.

Thus it is preferred that in a compound of formula (I):-

X is -OH or a lower alkoxy group containing from 1 to 4 carbon atoms or

X is R^5R^6N - wherein R^5 and R^6 are hydrogen or a lower alkyl group containing from 1 to 4 carbon atoms or

X is a group of formula IC'

$CH_2 - O -$

$CR^{18} - (R^{21})_d - O - (R^8O)_b - R^9 - NH_2$


$CH_2 - O - (R^{12}O)_c - R^{13} - NH_2$

wherein R^8O and $R^{12}O$ are independently ethoxy, or propoxy, in particular isopropoxy, or a random or block mixture thereof, R^9 and R^{13} are independently a linear or branched chain alkylene bridging group containing 2 or 3 carbon atoms and R^{21} is a linear or branched chain alkylene bridging group containing from 1 to 3 carbon atoms, R^{18} is hydrogen or lower alkyl containing from 1 to 4 carbon atoms and d is 0 or 1,

R^1 and R^2 are independently hydrogen or a lower alkyl group containing from 1 to 4 carbon atoms, R^3O is ethoxy, or propoxy, in particular isopropoxy, or a random or block mixture thereof, R^4 is a linear or branched chain alkylene bridging group containing 2 or 3 carbon atoms and a is from 1 to 400 or if X is of formula IC', the sum of a, b and c is from 3 to 400.

The compounds for use in the composition of the present invention are alkoxyated monoamines (X is of Formula IA), diamines (X is of Formula IB) or triamines (X is of Formula IC). Alkoxylation typically takes place to introduce ethoxy groups or propoxy groups, although butoxy groups may sometimes also be used. The propoxy group is preferably an isopropoxy group, $-OCH_2-CH(CH_3)-$. Mixed alkoxylation may also take place to introduce for example both ethoxy and propoxy or butoxy groups which may be present


as a mixture in either random or block arrangement. Thus each of $-(OR^3)_a-$, $-(OR^8)_b-$ and $-(OR^{12})_c-$ in the above formulae independently represent alkoxy groups, for example ethoxy and propoxy groups or a mixture thereof. Thus the definition of group $-(OR^3)_a$ for example includes a group



wherein a is represented by the sum of x and y. Other combinations of for example ethoxy propoxy and butyloxy groups in any desired order are similarly included in the definition of $-(OR^3)_a-$, $-(OR^8)_b-$ and $-(OR^{12})_c-$ respectively

The process of alkoxylation may produce a mixture of products having different
10 degrees of alkoxylation. Thus the values of a, b, c and a' represent an average degree of alkoxylation over the product as a whole. Preferred values of a, b, and c, respectively are from 1 to 100, for example from 1 to 50 and especially from 1 to 30. Thus when X in formula (I) is of formula IC, the sum of $a + b + c$ is preferably from 3 to 90. In the group $X'-(R^3O)_a-R^4-$ if present, a' is preferably from 0 to 50 and especially from 0 to 30.


15 An especially preferred alkoxyated monoamine (wherein X is of formula IA) for use in the present invention has the formula II wherein X is methoxy, $-(R^3O)_a$ represents a mixture of ethoxy and propoxy groups R^4 represents a propylene bridging group:-




The average degree of ethoxylation (x) may vary from 0 to about 45 or more preferably
20 from 0 to about 40, for example from 0 to about 20 and the average degree of propoxylation (y) may vary from 0 to about 90 and more preferably from about 1 to about 35, for example from about 2 to about 30, provided that x and y are not both 0 at the same time. Products are commercially available wherein x is about 1 and y is about 9; x is about 19 and y is about 3; x is about 6 and y is about 29 and x is about 32 and y is about 10. As specific
25 examples of commercially available products there may be mentioned JEFFAMINE M600 (JEFFAMINE is a trade mark of Huntsman Limited) having an approximate molecular weight of 600 and a propoxy to ethoxy ratio of 9 to 1, JEFFAMINE M1000 having an approximate molecular weight of 1000 and a propoxy to ethoxy ratio of 3 to 19, JEFFAMINE M2005 having an approximate molecular weight of 2000 and a propoxy to
30 ethoxy ratio of 29 to 6 and JEFFAMINE M2070 having an approximate molecular weight of 2000 and a propoxy to ethoxy ratio of 10 to 32.

Also included in the scope of the present invention is

(i) diethylene glycolamine wherein in Formula I, X is OH, (R³O) is ethoxy, a is 1 and R⁴ is an ethylene bridging group and R¹ and R² are hydrogen:-




5 (ii) a compound of formula



10 wherein in Formula I, X is -OH, R³ and R⁴ are isopropyl, R¹ and R² are X'-(R³O)_a-R⁴-, X' is -OH, R³ and R⁴ are ethyl and x, y and z respectively take any of the values defined for a or a' as the case may be and

(iii) a compound of formula




15 wherein in Formula I, X is R⁵R⁶N-, R³ and R⁴ are isopropyl, R¹, R², R⁵ and R⁶ are X'-(R³O)_a-R⁴-, X' is -OH, R³ and R⁴ are ethyl and v, w, x, y and z respectively take any of the values defined for a or a' as the case may be.

Also included in the scope of the present invention is

tris 2-(2-methoxyethoxy)ethylamine wherein in formula I, X is OCH₃, (R³O) is ethoxy, a is 1 and R⁴ is an ethylene bridging group and wherein R¹ and R² are each a group -R⁴-(OR³)_a-X' wherein (R³O) is ethoxy, a' is 1, R⁴ is an ethylene bridging group and X' is OCH₃.

25 An especially preferred propoxylated diamine (wherein X is of formula IB) for use in the present invention has the formula III wherein X is a group H₂N-, -(R³O)_a represents propoxy groups R⁴ represents a propoxy bridging group:-




30 wherein w is an average of from about 1 to about 80. Products are commercially available wherein w is an average of about 2.6, 5.6, 33.1 and 68. As Examples of commercially available products of formula III, there may be mentioned JEFFAMINE D230 having an approximate molecular weight of 230 and a value of w of about 2.6, JEFFAMINE D400 having an approximate molecular weight of 400 and a value of w of about 5.6,

35 JEFFAMINE D2000 having an approximate molecular weight of 2000 and a value of w of


about 33.1, and JEFFAMINE D4000 having an approximate molecular weight of 4000 and a value of w of about 68. An especially preferred value for w is from about 2 to about 35, for example from about 2 to about 30.

An alternative diamine (wherein X is of formula IB) having a mixture of ethoxylation and propoxylation for use in the present invention has the formula IV wherein X is a group H_2N- , $-(R^3O-)_s$ represents a mixture of ethoxy and propoxy groups R^1 and R^2 are hydrogen and R^4 represents a propylene bridging group:-



wherein the sum of $r + t$ (total propoxy content) is an average of from 1 to 20 propoxy groups, for example from about 1 to 10 propoxy units and s is an average of from 2 to 50 for example from 5 to 50 ethoxy units. Commercial products are available wherein the sum of $r + t$ is about 3.6 and s represents an average of about 9, or 15.5 respectively or wherein $r + t$ is about 6 and s represents an average of about 38.7. As examples of commercially available products of formula (IV) there may be mentioned JEFFAMINE ED 600 having an approximate molecular weight of 600 and an propoxy to ethoxy ratio of 3.6 to 9, JEFFAMINE ED 900 having an approximate molecular weight of 900 and an propoxy to ethoxy ratio of 3.6 to 15.5 and JEFFAMINE ED 2003 having an approximate molecular weight of 2000 and an propoxy to ethoxy ratio of 6.0 to 38.7. It is especially preferred that the value of $(r + t)$ is from about 2 to about 15 and s is from about 3 to about 20.

An especially preferred propoxylated triamine (wherein X is formula IC) has the formula (V)



wherein the sum of $d + e + f$ is an average of from about 5 to 90 and R^{22} is hydrogen or lower alkyl containing 1 to 4 carbon atoms such as methyl or ethyl, R^{21} is a linear or branched chain alkylene bridging group containing from 1 to 3 carbon atoms and d is 0 or 1. Products are available commercially wherein the sum of $d + e + f$ is about 5.6, about 50 and about 85 respectively. Thus as examples of commercially available products of formula (V) there may be mentioned JEFAMINE T403 having an approximate molecular weight of 440

and a value of $d + e + f$ (total propoxy content) of 5.6, JEFAMINE T3000 having an approximate molecular weight of 3000 and a value of $d + e + f$ (total propoxy content) of 50 and JEFAMINE T5000 having an approximate molecular weight of 5000 and a value of $d + e + f$ (total propoxy content) of 85.

5 We have found that the activity enhancing effect of the adjuvants in the composition of the present invention is particularly and surprisingly marked when the agrochemical active ingredient is water-soluble. The agrochemical for use in the composition of the present invention is preferably a water-soluble electrolyte such as an agrochemical salt and in particular a herbicidally active agrochemical salt for example a salt of glyphosate,
10 paraquat or fomesafen. Typical of the commonly available salts of glyphosate are the isopropylamine, trimethylsulphonium, sodium, potassium, ammonium, and ethanolamine salts. Paraquat is generally sold in the form of paraquat dichloride. Fomesafen is generally formulated as the sodium salt.

The proportion of adjuvant of formula (I) to agrochemical, for example glyphosate
15 (expressed as the acid) may vary within wide ranges depending on the desired level of activation. Typically the proportion of adjuvant of formula (I) to agrochemical will be from 1:40 to 3:1 for example from 1:20 to 3:1 by weight and especially from 1:10 to 1:1 by weight.

The compositions of the present invention may be used on their own but are
20 preferably used in the form of a composition containing a carrier comprising a solid or liquid diluent.

Compositions of the invention include both dilute compositions, which are ready for immediate use, and concentrated compositions, which require to be diluted before use, usually with water. The concentration of the composition will depend on the nature of the
25 active ingredient. Typically, and especially for example if the active ingredient is a herbicide, the compositions contain from 0.01% to 90% by weight of the active ingredient. Dilute compositions ready for use preferably contain from 0.01 to 2% of active ingredient, while concentrated compositions may contain from 20 to 90% of active ingredient, although from 20 to 70% is usually preferred.

30 The solid compositions may be in the form of granules, or dusting powders wherein the active ingredient and adjuvant are mixed with a finely divided solid diluent, e.g. kaolin, bentonite, kieselguhr, dolomite, calcium carbonate, talc, powdered magnesia, Fuller's earth

and gypsum. In some instances, and in particular when the active ingredient is glyphosate, inorganic salts such as ammonium sulphate may be used both as adjuvant and solid support. They may also be in the form of dispersible powders or grains, comprising a wetting agent to facilitate the dispersion of the powder or grains in liquid. Solid compositions in the form of a powder may be applied as foliar dusts.

The rate of application of the compositions of the invention will depend on a number of factors depending in particular on the nature of the active ingredient. When the active ingredient is a herbicide, such factors include, the compound chosen for use, the identity of the plants whose growth is to be inhibited, the formulations selected for use and whether the compound is to be applied for foliage or root uptake. As a general guide, however, an application rate of from 0.001 to 20 kilograms active ingredient per hectare is suitable while from 0.025 to 10 kilograms per hectare may be preferred.

The adjuvant of formula (I) or a mixture of different adjuvants of formula (I) may be used as the sole adjuvant to enhance the biological activity or physical properties of the agrochemical or may alternatively be used in conjunction with one or more additional adjuvants. The adjuvant of formula (I) may for example be combined with cationic surfactants, anionic surfactants, amphoteric surfactants or non-ionic surfactants. Such surfactants are well known in the art but as surfactants which are particularly well suited to be combined with an adjuvant of formula (I) there may be mentioned alkyl glycosides (mono and poly), alcohol ethoxylates, alkyl phenol ethoxylates, alkyl ester ethoxylates, sorbitan ester ethoxylates, siloxane ethoxylates, acetylenic diol ethoxylates, optionally alkoxyated tertiary or quaternary alkyl amines, optionally alkoxyated alkyl amine oxides, alkyl betaines optionally alkoxyated alkyl phosphate esters and sucrose alkyl esters. The adjuvant of formula (I) may also be combined with other activity-enhancing adjuvants, for example ammonium sulphate, urea or humectants, such as glycerol, polyethylene glycol, sorbitol, ethylene glycol, propylene glycol and lactate salts.

Certain of the adjuvants for use in the composition of the present invention have surprisingly low surface activity (high surface tension) as compared with conventional agrochemical adjuvants. These adjuvants with low surface activity generally exhibit a more favourable toxicological profile. Surface tension of the adjuvants for use in the present invention was measured as a 0.2% w/v solution in deionised water adjusted to pH 4 with hydrochloric acid at 20°C, and preferred adjuvants such as JEFFAMINE ED600 ED900 and

D400 have a surface tension measured by this method of greater than 50 mNm^{-1} as compared with AGRIMUL PG2067 alkylpolyglycoside which has a surface tension measured by this method of 28 mNm^{-1} . Adjuvants having a surface tension of less than 50 mNm^{-1} also give excellent results and JEFFAMINE D2000 for example having a surface tension measured by this method of 42 mNm^{-1} may even provide advantages on certain weed species as discussed below.

Compositions of the present invention generally exhibit excellent low-foaming properties when the composition is incorporated in the spray tank and during transportation and spraying.

When the active ingredient is a herbicide and in particular a water-soluble herbicide, adjuvants of formula (I) generally provide excellent and effective enhancement of activity in respect of most important weed species encountered in the field. Enhancement may however be less marked on certain "difficult" weed species. We have found that the use of a mixture of an adjuvant of formula (I) with a surfactant and in particular with an alkylpolyglycoside surfactant may provide excellent enhancement of activity both against normal weed species and against "difficult" species. Indeed in certain circumstances synergy may be observed and greater enhancement of control of "difficult" species may be obtained than when using either the adjuvant of formula (I) or the alkylpolyglycoside alone. The ratio of the adjuvant of formula (I) to the alkylglycoside is preferably from 1 : 40 to 4 : 1, for example 1 : 20 to 4 : 1, especially from 1 : 6 to 2 : 1 and most preferably about 1 to 1.

When the active ingredient is a herbicide, the invention provides, in a further aspect, a process for severely damaging or killing unwanted plants, which process comprises applying to the plants, or to the growth medium of the plants, a herbicidally effective amount of a composition as hereinbefore defined.

The invention is illustrated by the following Examples in which all parts and percentages are by weight unless otherwise stated.

Examples 1 to 10

Potassium glyphosate was applied at 500g glyphosate acid equivalent/ha to *Abutilon theophrasti* (ABUTH) plants grown in the glass-house. All treatments were made up in deionised water and applied using a tracksprayer with a 11002 nozzle at a spray application volume of 200l/ha. All treatments were replicated 4 times. After spraying the plants were

laid out in a glass-house and maintained at a temperature of 24°C by day and 19°C by night. The JEFFAMINE adjuvants were used in a proportion 0.2% w/v. A visual assessment of % control, where 0=unaffected and 100=complete kill, was carried out 16 days after treatment.

Example No	Treatment	% control
Comparison	No adjuvant	37
1	JEFFAMINE D400	93
2	JEFFAMINE D2000	76
3	JEFFAMINE T3000	79
4	JEFFAMINE ED600	89
5	JEFFAMINE ED900	85
6	JEFFAMINE ED2003	85
7	JEFFAMINE M600	85
8	JEFFAMINE M1000	81
9	JEFFAMINE M2005	70
10	JEFFAMINE M2070	74

5

EXAMPLES 11 TO 14

Potassium glyphosate was applied to *Abutilon theophrasti* (ABUTH), *Brassica napus* (BRSNS) and *Veronica persica* (VERPE) drilled as rows in the field, at a spray application volume of 200l/ha. The indicated adjuvant was included in the formulation at a total
 10 adjuvant concentration of 0.2% w/v. Application rates were 150, 300 and 450 g glyphosate acid/ha and each treatment was replicated 3 times. Data (mean of the three replicates and across all rates) is presented for a visual assessment of % control carried out at 26 days after application (DAA), where 0 = unaffected and 100% = complete kill.

AL2042 is an alkylpolyglycoside surfactant based on the same alkylpolyglycoside as
 15 AGRIMUL PG 2067. Both JEFFAMINE ED600 and JEFFAMINE D400 showed excellent adjuvant activity on representative species ABUTH and VERPE. On the "difficult" species BRSNS, the combination of the JEFFAMINE adjuvant and the alkylpolyglycoside showed greater activity than either the JEFFAMINE adjuvant or the alkylpolyglycoside on its own.

Example No	Total adjuvant = 0.2%	ABUTH	BRSNS	VERPE
11	JEFFAMINE ED600	70.0	64.7	48.8
12	JEFFAMINE D400	64.9	62.2	47.2
13	JEFFAMINE ED600 and Agrimul PG2067 (0.1% + 0.1%)	57.8	78.9	44.4
14	JEFFAMINE D400 and Agrimul PG2067 (0.1% + 0.1%)	57.7	77.1	44.7
Comparison	AL2042	45.8	70.3	39.7

EXAMPLE 15

This Example illustrates the use of a salt of ethoxy (5 moles of EO) isotridecyl phosphate ester in the acid form (CRODAFOS T5A) with JEFFAMINE ED600.

- 5 CRODAFOS T5A is a mixture of mono and di esters. CRODAFOS is a tradename of Croda

Potassium glyphosate, in combination with the adjuvants specified below, was applied to *Ipomoea hederacea* (IPOHE) at 300 g glyphosate acid equivalent/ha. All treatments were made up in deionised water and applied using a tracksprayer with a 11002 nozzle at a spray application volume of 200l/ha. All treatments were replicated 4 times.

- 10 After spraying the plants were laid out in a glass-house and maintained at a temperature of 24°C by day and 19°C by night. A visual assessment of % control was carried out 16 days after treatment

Treatment	IPOHE - % control
No adjuvant	64
AL2042	71
CRODAFOS T5A, JEFFAMINE ED600 salt	86
CRODAFOS T5A, potassium salt	79

15

The AL2042 was applied at 0.2% w/v and the CRODAFOS T5A salts were applied at 0.2% w/v phosphate ester acid equivalent. The Crodafos T5A potassium salt was prepared by neutralising the phosphate ester in its acid form with potassium hydroxide to pH6. The CRODAFOS T5A JEFFAMINE ED600 salt was prepared by neutralising the phosphate ester in its acid form with JEFFAMINE ED600 to pH6.

20

EXAMPLE 16

This Example illustrates the use of salts of glyphosate acid and the adjuvants of Formula (I).

- 5 Glyphosate trimesium and the glyphosate salts of the indicated adjuvants of formula (I) were applied to *Sorghum halepense* (SORHA) at 500, 1000, 2000 and 4000 g glyphosate acid equivalent/ha. All treatments were made up in deionised water and applied using a tracksprayer with a 11002 nozzle at a spray application volume of 200l/ha. All treatments were replicated 3 times. After spraying the plants were laid out in a glass-house and
- 10 maintained at a temperature of 24°C by day and 19°C by night. A visual assessment of % control was carried out 22 days after treatment and calculated doses for 90% control (ED90) were derived. AL2042 alkylpolyglycoside was used at 0.25% w/v as additional surfactant with all treatments. It will be appreciated that a low ED90 value indicates high activity.

Glyphosate salt	Adjuvant	SORHA - ED90 gae/ha
Glyphosate trimesium	AL2042	1410
Glyphosate JEFFAMINE D400	AL2042	821
Glyphosate JEFFAMINE M600	AL2042	493

15


EXAMPLE 17

- Paraquat dichloride, in combination with the adjuvants specified below, was applied to *Abutilon theophrasti* (ABUTH) at four rates of paraquat (15, 45, 135 and 270 g/ha paraquat ion). All treatments were made up in deionised water and applied using a
- 20 tracksprayer with a 11002 nozzle at a spray application volume of 200l/ha. All treatments were replicated 3 times. After spraying the plants were laid out in a glass-house and maintained at a temperature of 22°C by day and 18°C by night. A visual assessment of % control was carried out 18 days after treatment and calculated doses for 90% control (ED90 - g paraquat ion per litre) were calculated. The adjuvants were applied at 0.1% w/v.
- 25 ED90 g/ha 18DAT

Treatment (0.1% adjuvant)	ABUTH	SIDSP
Control - No adjuvant	325	(>270)
JEFFAMINE ED600	246	182

Claims

1. An agrochemical composition comprising an agrochemical active ingredient and an adjuvant of formula (I) and salts thereof



wherein R^1 and R^2 are independently hydrogen or a lower alkyl group or a group $X'-(R^3O)_a-R^4$ - wherein R^3 and R^4 respectively may take any of the values of R^3 and R^4 as hereinafter defined and wherein X' is -OH or a lower alkoxy group containing from 1 to 6 carbon atoms and a is from 0 to 400

R^3O is an ethoxy, propoxy or butoxy group or a random or block mixture thereof;

R^4 is a linear or branched chain alkylene bridging group containing from 1 to 4 carbon atoms, X is

(IA) -OH or a lower alkoxy group containing from 1 to 6 carbon atoms or X is

(IB) a group R^5R^6N - or $R^5R^6N-R^7$ -

wherein R^5 and R^6 are independently hydrogen or a lower alkyl group or a group $X'-(R^3O)_a-R^4$ - as hereinbefore defined wherein X' is -OH or a lower alkoxy group containing from 1 to 6 carbon atoms,

and R^7 is a linear or branched chain alkylene bridging group containing from 1 to 4 carbon atoms

or X is

(IC) a group

$CR^{16}R^{17}-O-$

$|$
 $CR^{18}-(R^{21})_d-O-(R^8O)_b-R^9-NR^{10}R^{11}$

$|$
 $CR^{19}R^{20}-O-(R^{12}O)_c-R^{13}-NR^{14}R^{15}$

wherein R^{10} , R^{11} , R^{14} and R^{15} are independently hydrogen or a lower alkyl group or a group $X'-(R^3O)_a-R^4$ - as hereinbefore defined wherein X' is -OH or a lower alkoxy group containing from 1 to 6 carbon atoms,

and R^8 and R^{12} are independently alkyl groups containing from 1 to 4 carbon atoms and R^9 , R^{13} and R^{21} are independently a linear or branched chain alkylene bridging group containing from 1 to 6 carbon atoms, R^{16} , R^{17} , R^{18} , R^{19} and R^{20} are independently hydrogen or a lower alkyl group and d is 0 or 1

and wherein a, is from 1 to 400 or if X is of formula IC, the sum of a, b and c is from 3 to 400.

2. A composition according to claim 1 wherein

X is -OH or a lower alkoxy group containing from 1 to 4 carbon atoms or

X is R^5R^6N - wherein R^5 and R^6 are hydrogen or a lower alkyl group containing from 1 to 4 carbon atoms or

X is a group of formula IC'

$CH_2 - O -$

$|$
 $CR^{18} - (R^{21})_d - O - (R^8O)_b - R^9 - NH_2$


$|$
 $CH_2 - O - (R^{12}O)_c - R^{13} - NH_2$

wherein R^8O and $R^{12}O$ are independently ethoxy, or propoxy or a random or block mixture thereof, R^9 and R^{13} are independently a linear or branched chain alkylene bridging group containing 2 or 3 carbon atoms, R^{21} is a linear or branched chain alkylene bridging group containing from 1 to 3 carbon atoms, R^{18} is hydrogen or lower alkyl containing from 1 to 4 carbon atoms, d is 0 or 1,

R^1 and R^2 are independently hydrogen or a lower alkyl group containing from 1 to 4 carbon atoms, R^3O is ethoxy, or propoxy or a random or block mixture thereof, R^4 is a linear or branched chain alkylene bridging group containing 2 or 3 carbon atoms and a is from 1 to 400 or if X is of formula IC', the sum of a, b and c is from 3 to 400.

3. A composition according to claim 1 or 2 wherein a is from 1 to 50 or if X is of formula IC or IC' respectively, the sum of a, b and c is from 3 to 90 and wherein in the group $X' - (R^3O)_a - R^4$ - if present, a' is from 0 to 50.


4. A composition according to claim 1 wherein the adjuvant of formula (I) is an alkoxyated monoamine, having the formula II



wherein the average degree of ethoxylation (x) is from 0 to about 45 and the average degree of propoxylation (y) is from 0 to about 90 provided that x and y are not both 0 at the same time.

5. A composition according to claim 4 wherein the value of x is from 0 to about 20 and of y is from about 2 to about 30.


6. A composition according to claim 1 wherein the compound of formula (I) is a diamine having a formula (II)



wherein w is an average of from about 1 to about 80.

7. A composition according to claim 6 wherein w is an average of from about 2 to about 35


8. A composition according to claim 1 wherein the compound of formula (I) is a diamine having a formula (IV)



wherein the sum of r + t is an average of from 1 to 20 propoxy groups and s is an average of from 2 to 50 ethoxy units.

9. A composition according to claim 8 wherein the sum of r + t is from about 2 to about 15 and s is from about 3 to about 20.

10. A composition according to claim 1 wherein the compound of formula (I) is a propoxylated triamine having the formula (V)



wherein the sum of d + e + f is an average of from about 5 to 90, R²² is hydrogen or lower alkyl containing 1 to 4 carbon atoms, R²¹ is a linear or branched chain alkylene bridging group containing 1 to 3 carbon atoms and d is 0 or 1.

11. A composition according to any of the preceding claims wherein the proportion of adjuvant of formula (I) to the agrochemical is from 1:20 to 3:1 by weight.

12. A composition according to any of the preceding claims wherein the adjuvant of formula (I) in claim 1 is used in combination with an alkylpolyglycoside surfactant.

13. A composition according to claim 12 wherein the ratio of the adjuvant of formula (I) to the alkylpolyglycoside surfactant is from 1 : 40 to 4 : 1.
14. A composition according to any of the preceding claims wherein the agrochemical active ingredient is a salt of glyphosate, a salt of fomesafen or a paraquat salt.
- 5 15. A composition according to any of claims 1 to 13 wherein the adjuvant of formula (I) in claim 1 is used in the form of a salt with glyphosate or a salt with an acidic surfactant.
16. A process of severely damaging or killing unwanted plants which process comprises applying to the plants or to the growth medium of the plants, a herbicidally effective
10 amount of a composition as claimed in claim 14.
17. A composition according to any of claims 1 to 15 which is an aqueous composition containing from 0.01% to 90% by weight of the active ingredient.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/GB 00/01062

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A01N25/30 A01N57/20

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP 0 375 624 A (SANDOZ AG) 27 June 1990 (1990-06-27) page 2, line 1 -page 3, line 8 compositions IVa, IVb, IVc, IVd example 3	1-3, 6, 11, 14-17
X	WO 96 32839 A (MONSANTO COMPANY) 24 October 1996 (1996-10-24) page 6, line 7 - line 28 page 14, line 23 - line 30	1-3, 11-17
X	WO 97 32476 A (KAO CORPORATION) 12 September 1997 (1997-09-12) page 2, line 23 -page 5, line 18 page 10, line 17 -page 11, line 16 page 17, line 14 -page 18, line 5	1-3, 11-17
	-/--	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

22 May 2000

Date of mailing of the international search report

07/06/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Fort, M

INTERNATIONAL SEARCH REPORT

Inter. nal Application No

PCT/GB 00/01062

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 5 668 085 A (JAMES CARNEGIE FORBES) 16 September 1997 (1997-09-16) the whole document	1-3,11, 14-17
X	WYRILL J B ET AL: "GLYPHOSATE TOXICITY TO COMMON MILKWEED AND HEMP DOGBANE AS INFLUENCED BY SURFACTANTS" WEED SCIENCE,US,WEED SCIENCE SOCIETY OF AMERICA, CHAMPAIGN, IL, vol. 25, no. 3, 1 May 1977 (1977-05-01), pages 275-287, XP002034447 ISSN: 0043-1745 page 277 -page 278	1-3,11, 14-17

INTERNATIONAL SEARCH REPORT

...formation on patent family members

International Application No

PCT/GB 00/01062

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 375624	A	27-06-1990	
		US 5035738 A	30-07-1991
		AT 118671 T	15-03-1995
		AU 629278 B	01-10-1992
		AU 4717189 A	28-06-1990
		BR 8906695 A	11-09-1990
		CA 2006114 A	23-06-1990
		CN 1043601 A,B	11-07-1990
		DE 68921317 D	30-03-1995
		DK 656989 A	24-06-1990
		EG 19012 A	30-06-1994
		ES 2070927 T	16-06-1995
		GR 3015295 T	30-06-1995
		HU 52685 A,B	28-08-1990
		JP 2221204 A	04-09-1990
		JP 2828291 B	25-11-1998
		NZ 231966 A	25-11-1992
		PT 92673 A,B	08-01-1991
		RU 2093505 C	20-10-1997
		TR 24048 A	01-03-1991
		ZA 8909864 A	28-08-1991
WO 9632839	A	24-10-1996	
		US 5750468 A	12-05-1998
		AT 187602 T	15-01-2000
		AU 691425 B	14-05-1998
		AU 5325196 A	07-11-1996
		BR 9608102 A	20-07-1999
		CA 2214376 A	24-10-1996
		CZ 9703327 A	18-02-1998
		DE 69605660 D	20-01-2000
		EP 0820231 A	28-01-1998
		ES 2142055 T	01-04-2000
		HU 9801372 A	28-09-1998
		JP 3015107 B	06-03-2000
		JP 10513478 T	22-12-1998
		NZ 305278 A	25-02-1999
		PL 322682 A	16-02-1998
		ZA 9602816 A	02-10-1996
WO 9732476	A	12-09-1997	
		AU 713956 B	16-12-1999
		AU 2231597 A	22-09-1997
		BR 9702106 A	12-01-1999
		CN 1190327 A	12-08-1998
		EP 0824315 A	25-02-1998
		JP 9295901 A	18-11-1997
		NZ 328990 A	29-06-1999
		US 5948421 A	07-09-1999
US 5668085	A	16-09-1997	NONE

